

Angiotensin II receptor antagonists: the development of the pathophysiological and clinical research

Ettore Ambrosioni, Stefano Bacchelli

Department of Internal Medicine D. Campanacci, University of Bologna, Bologna, Italy

(Ital Heart J 2000; 1 (Suppl 3): S12-S14)

Address:

Dr. Stefano Bacchelli

Dipartimento di Medicina
Clinica e Biotecnologia
Applicata "D. Campanacci"
Università degli Studi
Polclinico S. Orsola-
Malpighi
Via Massarenti, 9
40138 Bologna

Neurohormonal activation is one of the hallmarks of chronic heart failure and inhibition of the neurohormonal systems, in particular the sympathetic nervous system and the renin-angiotensin-aldosterone system (RAAS), have become important targets in heart failure. Indeed, the favorable effect of angiotensin-converting enzyme (ACE) inhibitors in heart failure has largely been attributed to the blockade of angiotensin II production, the target hormone of the RAAS, with favorable effects on the clinical course, including prolonged survival¹⁻⁴. Angiotensin II also contributes to the generalized (and mostly deleterious) activation of other neurohormonal systems both within the heart and throughout the body by promoting the release of agents such as norepinephrine, endothelin and aldosterone. The importance of aldosterone in heart failure has been overlooked in recent years because ACE-inhibitor-related reductions in angiotensin were thought to eliminate aldosterone production. Such suppression of circulating aldosterone, however, is transient, as exemplified by the term "escape" used to describe the phenomenon⁵. Indeed, the blockade of aldosterone receptors by spironolactone, in addition to standard therapy comprehensive of ACE-inhibitors, substantially reduces the risk of both morbidity and death among patients with severe heart failure⁶.

In addition, many of the benefits of the blockade of the RAAS in the field of heart failure appear to be related to the ability of the drugs to inhibit further cardiac remodeling⁷. What is not clear is the mechanism by which ACE-inhibitors lead to favorable changes in cardiac structure and function. Other than to block the production of an-

giotensin II through the converting-enzyme pathway, ACE-inhibitors also block the breakdown of bradykinin that reduces vasomotor tone by enhancing the release of vasodilator substances from the vascular endothelium, such as nitric oxide. It has also been shown to have antigrowth properties, which, at least in some animal experiments, account for the favorable effects of ACE-inhibitors on cardiac remodeling^{7,8}.

In addition to ACE-inhibitors and aldosterone-receptor blockers, there are now alternative pharmacological approaches to blocking the RAAS.

Development of a novel mechanism for interrupting the renin-angiotensin-aldosterone system

Selective blockade of the angiotensin II AT₁ receptor represents a novel mechanism for interrupting the RAAS without altering the potential benefits of AT₂ receptor stimulation⁹⁻¹¹.

Selective blockade of the angiotensin AT₁ receptor has several possible advantages over inhibition of ACE. Urata et al.¹²⁻¹⁴ have shown that there are a number of ACE-independent pathways for angiotensin II formation in human tissues and cardiovascular diseases, so that the conversion of angiotensin I to angiotensin II may involve a chymase-dependent pathway in addition to ACE. Moreover, activation of angiotensin II-forming chymase has been observed in various disease models such as cardiomyopathy, chronic mitral regurgitation as well as in the failing human heart¹¹. This so-called "escape" from suppression of the RAAS during ACE-

inhibition is so often encountered in circumstances of increased activity, but its clinical relevance is largely unknown¹⁵. Very recently, Roig et al.¹⁶ have found that 50% of patients had increased angiotensin II levels, despite high-dose ACE-inhibitors, and that their prognosis was worse compared to patients in whom angiotensin II remained suppressed.

Another potential advantage of selective blockade of the angiotensin AT₁ receptor is the enhanced stimulation of the angiotensin AT₂ receptor. Evidence suggests that stimulation of the AT₂ receptor may have beneficial effects on the vasculature, although the consequences of activation of these receptors are only beginning to be understood. Several studies have shown that stimulation of angiotensin AT₂ receptors inhibits cell proliferation and that angiotensin II-induced cardiomyocyte hypertrophy is increased by selective blockade of the AT₂ receptor¹⁷⁻¹⁹. Interestingly, it has recently been suggested that angiotensin II receptor blockers, in addition to blockade of the angiotensin II AT₁ receptor, might stimulate bradykinin through the unprotected AT₂ receptor²⁰. Although these findings will need further confirmation, ACE-inhibitors decrease stimulation of the AT₂ receptor, while angiotensin II receptor blockers increase stimulation of this receptor because of the increase in circulating levels of angiotensin II that interact with other unblocked angiotensin II receptors.

Taken together, these findings suggest that the vascular protective effects observed with AT₁ receptor antagonists may be at least partially caused by an unopposed antigrowth effect of AT₂ receptor stimulation. In contrast, such a response would not be obtained with ACE-inhibitors, since these drugs equally antagonize AT₁ and AT₂ receptors by inhibiting the generation of angiotensin II¹¹.

Furthermore, the blockade of the angiotensin AT₁ receptor prevents some of the difficulties associated with ACE-inhibitor treatment²¹, that have a number of side effects including dry cough, angioedema, and urticaria. There is some evidence suggesting that the side effects associated with ACE-inhibitors are not produced by inhibition of angiotensin II synthesis but rather inhibition of kininase 2, which leads to a reduction in bradykinin metabolism.

Selective blockade of AT₁ receptors, therefore, not only represents an alternative mechanism for interrupting the RAAS but potentially offers several therapeutic advantages compared with ACE-inhibition, and the difference in mode of action theoretically favors add-on therapy and raises the intriguing possibility that combination therapy with both classes of drugs could offer benefits beyond those seen with either of the agents alone⁷.

This hypothesis was confirmed in a recent double-blind study in a small group of patients with heart failure, in which clinical conditions improved when an AT₁ receptor blocker was added to maximal ACE-inhibition²². The combination of candesartan and enalapril pre-

vented an increase in left ventricular volume that occurred with either of the drugs alone and appeared to have favorable effects on the neurohormonal profile of these patients, despite a trend toward an increased number of events in patients treated with candesartan²³.

However, the recently published ELITE (Evaluation of Losartan in the Elderly) II study has not demonstrated a significant difference in terms of all-cause mortality in patients with heart failure treated with losartan in comparison with an ACE-inhibitor, despite a better tolerability profile of the AT₁ receptor blocker²⁴. In the near future other data from ongoing clinical studies will be available. One of these, the Val-HeFT (Valsartan Heart Failure Trial), is evaluating the effect of AT₁ receptor blockade used as add-on therapy in patients already receiving ACE-inhibitors²⁵. Another one, the CHARM (Candesartan in Heart Failure Assessment of Reduction in Mortality and Morbidity), is investigating the role of AT₁ blockade in patients with left ventricular systolic dysfunction and intolerant to ACE-inhibition, or in addition to ACE-inhibitors; furthermore, the third arm of this study is studying the effects of candesartan in heart failure patients with diastolic dysfunction (left ventricular ejection fraction > 0.40)²⁵.

The results of these large trials are being extensively awaited and will provide important information about the exact place of AT₁ receptor blockers in patients with heart failure. Until such time, however, ACE-inhibitors remain the cornerstone of the treatment in heart failure, despite their limitations, since their value in these patients is beyond doubt¹⁵.

References

1. The CONSENSUS Trial Study Group. Effects of enalapril on mortality in severe congestive heart failure: results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). *N Engl J Med* 1987; 316: 1429-35.
2. Cohn JN, Johnson G, Ziesche S, et al. A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med* 1991; 325: 303-10.
3. The SOLVD Investigators. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991; 325: 293-302.
4. The SOLVD Investigators. Effect of enalapril on mortality and the development of heart failure in asymptomatic patients with reduced left ventricular ejection fraction. *N Engl J Med* 1992; 327: 685-91.
5. Weber KT. Aldosterone and spironolactone in heart failure. *N Engl J Med* 1999; 341: 753-5.
6. Pitt B, Zannad F, Remme WJ, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. *N Engl J Med* 1999; 341: 709-17.
7. Greenberg BH. Role of angiotensin receptor blockers in heart failure. Not yet RESOLVD. *Circulation* 1999; 100: 1032-4.
8. McDonald KM, Garr M, Carlyle PF, et al. Relative effects of alpha 1-adrenoceptor blockade, converting enzyme inhibitor therapy, and angiotensin II subtype 1 receptor blockade on ventricular remodeling in the dog. *Circulation* 1994; 90: 3034-46.

9. Ambrosioni E, Bacchelli S. Farmacologia clinica degli antagonisti dell'angiotensina II. *Cardiologia* 1994; 39 (Suppl 1 al n 12): 401-4.
10. Horiuchi M, Akishita M, Dzau VJ. Recent progress in angiotensin II type 2 receptor research in the cardiovascular system. *Hypertension* 1999; 33: 613-21.
11. Brooks DP, Ohlstein EH, Ruffolo RR. Pharmacology of eprosartan, an angiotensin II receptor antagonist: exploring hypotheses from clinical data. *Am Heart J* 1999; 138: S246-S251.
12. Urata H, Nishimura H, Ganten D, et al. Angiotensin converting enzyme-independent pathways of angiotensin II formation in human tissues and cardiovascular diseases. *Blood Press* 1996; 5: 22-8.
13. Urata H, Kinoshita A, Misono KS, Bumpus FM, Husain A. Identification of a highly specific chymase as the major angiotensin II-forming enzyme in the human heart. *J Biol Chem* 1990; 265: 22348-57.
14. Urata H, Nishimura H, Ganten D. Chymase-dependent angiotensin II forming system in humans. *Am J Hypertens* 1996; 9: 277-84.
15. Van Veldhuisen DJ, Voors AA. Blockade of the renin angiotensin system in heart failure: the potential place of angiotensin II receptor blockers. *Eur Heart J* 2000; 21: 14-6.
16. Roig E, Perez-Villa F, Morales M, et al. Clinical implications of increased plasma angiotensin II despite ACE inhibitors therapy in patients with congestive heart failure. *Eur Heart J* 1999; 20: 53-7.
17. Stoll M, Steckelings UM, Paul M, et al. The angiotensin AT₂ receptor mediates inhibition of cell proliferation in coronary endothelial cells. *J Clin Invest* 1995; 95: 651-7.
18. Tsuzuki S, Mataba T, Eguchi S, et al. Angiotensin II type 2 receptor inhibits cell proliferation and activates tyrosine phosphatase. *Hypertension* 1996; 28: 916-8.
19. Booz GW, Baker KM. Role of type 1 and type 2 angiotensin receptor in angiotensin II-induced cardiomyocyte hypertrophy. *Hypertension* 1996; 28: 635-40.
20. Gohlke P, Pees C, Unger T. AT₂ receptor stimulation increases aortic cyclic GMP in SHRSP by a kinin-dependent mechanism. *Hypertension* 1998; 31: 349-55.
21. Mann RD, Mackay F, Pearce G, Freemantle S, Wilton LV. Losartan: a study of pharmacovigilance data on 14 522 patients. *J Hum Hypertens* 1999; 13: 551-7.
22. Hamroff G, Katz SD, Mancini D, et al. Addition of angiotensin II receptor blockade to maximal angiotensin-converting enzyme inhibition improves exercise capacity in patients with severe congestive heart failure. *Circulation* 1999; 99: 990-2.
23. McKelvie RS, Yusuf S, Pericak D, et al. Comparison of candesartan, enalapril, and their combination in congestive heart failure: Randomized Evaluation of Strategies for Left Ventricular Dysfunction (RESOLVD) Pilot Study. *Circulation* 1999; 100: 1056-64.
24. Pitt B, Poole-Wilson PA, Segal R, et al. Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomised trial - the Losartan Heart Failure Survival Study ELITE II. *Lancet* 2000; 355: 1582-7.
25. Willenheimer R, Dahlöf B, Rydberg E, Erhardt L. AT₁-receptor blockers in hypertension and heart failure: clinical experience and future directions. *Eur Heart J* 1999; 20: 997-1008.